

ORIGINAL PAPER

Critical care admission trends and outcomes in individuals with bronchiectasis in the UK

Vidya Navaratnam^{1,2}, Colin R. Muirhead^{2,3}, Richard B. Hubbard^{1,2} and Anthony De Soyza^{2,4}

From the ¹Division of Epidemiology and Public Health, University of Nottingham, Nottingham, UK, ²BronchUK, a MRC Funded Partnership in Bronchiectasis, ³Institute of Health and Society, Newcastle University, Newcastle upon Tyne, UK and ⁴Institute of Cellular Medicine, Newcastle University & Sir William Leech Centre, Newcastle upon Tyne, UK

Address correspondence to Dr Vidya Navaratnam, Nottingham Respiratory Research Unit, Clinical Sciences Building, Nottingham City Hospital, Hucknall Road, Nottingham NG5 1PB, UK. email: vidya.navaratnam@nottingham.ac.uk

Summary

Background: There are limited data on admission trends and outcomes of individuals with bronchiectasis admitted to intensive care (ICU). Using national critical care data, we analysed admissions to ICU and estimated outcomes in terms of mortality in individuals with bronchiectasis and chronic obstructive pulmonary disease (COPD) admitted to ICU.

Methods: Using data from the Intensive Care National Audit and Research Centre, admissions from bronchiectasis and COPD from 1 January 2009 to 31 December 2013 were extracted. Crude admission rates for bronchiectasis and COPD were calculated and Poisson regression was used to estimate unadjusted annual admission rate ratios. We investigated changes to length of stay on ICU, ICU mortality and in-hospital mortality during the study period. We also compared mortality rates in people with bronchiectasis and COPD aged 70 or above.

Results: We found an annual increase of 8% (95% Confidence Interval [CI] 2–15) in the number of ICU admissions from bronchiectasis, whilst the yearly increase in ICU admissions from COPD was 1% (95% CI 0.3–2). ICU and in-hospital mortality was higher in individuals with bronchiectasis compared with those with COPD, especially in people aged 70 years or above.

Conclusion: Admission to ICU in people with bronchiectasis are uncommon, but are increasing in frequency over time, and carries a substantial mortality rate. This needs to be considered allocating health care resources and planning respiratory services.

Introduction

Bronchiectasis is a chronic lung disease characterized by repeated episodes of infection and chronic inflammation.¹ Recent studies suggest that incidence,² hospital admissions³ and mortality⁴ from bronchiectasis in the UK are increasing, but there are no data on admission rates to intensive care (ICU). There are also limited data on outcomes of individuals with

bronchiectasis admitted to ICU in terms of survival. These data are important to help health care planning and form guidance for thresholds for admission to critical care.

The aim of this study was to investigate trends in bronchiectasis admissions to ICU in the UK and estimate outcomes in patients with bronchiectasis admitted to ICU. To benchmark these data, we compared them with data on patients with chronic

Received: 3 August 2015; Revised (in revised form): 15 October 2015

© The Author 2015. Published by Oxford University Press on behalf of the Association of Physicians.
All rights reserved. For Permissions, please email: journals.permissions@oup.com

obstructive pulmonary disease (COPD), a common chronic respiratory disease in the UK. We also estimated financial impact of bronchiectasis admissions to ICU on secondary care services.

Methods

Data source

We used data from the Intensive Care National Audit and Research Centre (ICNARC), an electronic database of patient outcomes from adult general critical care units across England, Wales and Northern Ireland.⁵ 99% of adult critical care units contribute data to ICNARC, which includes information from 1.5 million individuals. Data from ICU admissions from adults over the age of 18 with bronchiectasis and COPD from 1 January 2009 to 31 December 2013 were extracted.

Definition of outcomes

Our definition of bronchiectasis admissions to ICU included patients whose main reason for admission to ICU was exacerbation of bronchiectasis and excluded individuals with cystic fibrosis. COPD admissions were defined as admissions whose main reason for admission was either COPD with acute lower respiratory infection; or COPD with acute exacerbation, cause unspecified. Patients who had COPD-bronchiectasis overlap⁶ were excluded from these analyses ($n=36$). The ICNARC Coding Method is a hierarchical method specifically designed for coding reasons for admission to critical care. Primary reason for admission is mandated; however, secondary reason for admission is optional, unless the patient was admitted following surgery.

ICU mortality was defined as the status at discharge from the ICU, whilst in-hospital mortality was defined as the status on discharge from hospital. We also extracted demographic information, intubation rates, length of stay on ICU (duration from date of admission to date of discharge from ICU or date of death), total hospital length of stay (date of admission to date of discharge from hospital or date of death), as well as mean Acute Physiology and Chronic Health Evaluation II (APACHE II) scores.

Statistical analysis

Only summary data grouped by calendar year were available for these analyses. We calculated crude admission rates for bronchiectasis and COPD for each year by dividing the number of admissions from bronchiectasis or COPD by the total number of admissions to ICU. Poisson regression modelling was used to estimate crude annual admission rate ratios. We calculated the cost of critical admissions from bronchiectasis for each year by multiplying the cost of a critical care bed day in the NHS with single organ support for the 2012–2013 financial year (www.dh.gov.uk) by the median length of stay on ICU and number of admissions for each calendar year. We developed an unadjusted Poisson regression model to predict future critical care admissions from bronchiectasis for the year 2020 and estimated cost of critical care admissions for that year without adjusting for inflation costs or changes to market forces. We investigated changes in mean age at admission, mean APACHE II scores, rates of intubation, ICU and in-hospital mortality over the study period in people with bronchiectasis and COPD. Finally, in separate analyses we compared these features between patients with COPD and bronchiectasis who were 70 years or older.

Two-sided likelihood ratio, unpaired t tests, Chi-squared and Wilcoxon rank sum tests were used for hypothesis testing.

Stata v12 (College Station, TX) was used for all statistical analyses.

Results

There were 614 352 admissions to ICU across 219 critical care units between 2009 and 2013, of which 536 (0.1%) were due to bronchiectasis and 19 754 (3.2%) due to COPD. Admissions from bronchiectasis increased from 74 in 2009 to 121 in 2013, equating to a crude annual increase of 8% (rate ratio [RR] 1.08; 95% Confidence Interval [CI] 1.02–1.15; $P=0.01$). Individuals with bronchiectasis were more frequently female (57.5%) and the mean age at admission increased from 56.6 (SD 18) to 65.8 years (SD 15.2) during the study period (linear regression $P=0.042$). However, rates of intubation (47.9 vs. 44.6%; $P=0.65$) and mean APACHE II acute physiology scores remained similar (12.7 vs. 13.5; $P=0.17$) during this time. ICU mortality did not change (27.0 vs. 28.9%; $P=0.83$), neither did in-hospital mortality (35.7 vs. 33.9%; $P=0.86$) (see Table 1). There was also no change in median length of stay in ICU ($P=0.21$) or median length of stay in hospital ($P=0.36$) (see Table 2). During the 5-year study period, the estimated financial burden on the National Health Service (NHS) in terms of critical care bed days from bronchiectasis increased marginally from £189 144 to £298 967 per year (see Table 2). Using an unadjusted Poisson regression model, we estimate that by 2020, there will be close to 220 critical care admissions per year from bronchiectasis, which will cost the health service almost £500 000 annually in critical care bed day costs.

In contrast, there were 3126 admissions due to COPD in 2009, which rose to 4652 in 2013. The unadjusted year-on-year increase in COPD admissions was 1% (RR 1.010 95% CI 1.003–1.023; $P=0.012$). The mean age at admission remained static (67.5 years [SD 10.6] in 2009 vs. 67.9 years [SD 10.6] in 2013; $P=0.16$), but ICU mortality decreased during the study period (20.5 vs. 18.0%; $P=0.02$), as did in-hospital mortality (30.7 vs. 26.7%; $P=0.005$) (see Table 1). The proportion of individuals with COPD who were intubated also decreased during this time (53.7 vs. 44.2%; $P<0.001$).

In separate analyses of 219 people with bronchiectasis aged 70 or above, we found that ICU mortality was higher compared with that amongst those aged under 70 (31.1 vs. 18.3%; $P<0.001$) despite having similar mean APACHE II acute physiology scores. Furthermore, in-hospital mortality in individuals with bronchiectasis aged 70 or above was almost double compared with individuals with bronchiectasis under 70 (47.4 vs. 24.6%; $P<0.001$). In comparison to people with COPD aged 70 or above, we found that patients with bronchiectasis aged 70 or above had substantially higher rates of death whilst on ICU (23.1 vs. 31.1%; $P=0.03$) and during hospital admission (36.4 vs. 46.6%; $P=0.04$) (see Table 3). Individuals with bronchiectasis over the age of 70 were also less likely to be intubated compared with those <70 years. However, the proportions of individuals over the age of 70 with bronchiectasis and COPD who were intubated were similar (41.6 vs. 42.2%; $P=0.90$) (see Table 3).

Discussion

To the best of the authors' knowledge, this is the first study to use national data to determine trends in ICU admissions amongst individuals with bronchiectasis. We found that the number of patients with bronchiectasis admitted to ICU is increasing. The mean age at admission increased during our study period but there was no change to length of stay on ICU.

Table 1. Crude admission rates, Poisson regression modeling of admissions, mortality and median ICU length of stay in people with Bronchiectasis and COPD

Bronchiectasis	Number of admissions	Person years	Crude admission rate per 100 000 person years (95% CI)	Crude admissions rate ratio (95% CI)	Number of ICU deaths (%)	Number of hospital deaths (%)
Year						
2009	74	97 457	75.9 (60.5–95.4)	1.00	20 (27.0)	25 (35.7)
2010	78	114 207	68.3 (54.7–85.3)	0.90 (0.65–1.24)	18 (23.1)	31 (41.3)
2011	108	128 659	83.9 (69.5–101.4)	1.11 (0.82–1.49)	22 (20.4)	32 (31.4)
2012	155	137 062	113.1 (96.6–132.7)	1.49 (1.13–1.96)	31 (20.0)	48 (31.8)
2013	121	136 877	88.4 (74.0–105.6)	1.16 (0.87–1.55)*P for trend = 0.022	35 (28.9)	40 (33.9)
COPD						
Year						
2009	3126	97 457	3204.6 (3094.2–3318.9)	1.00	642 (20.5)	941 (30.7)
2010	3675	114 207	3217.8 (3115.5–3223.6)	1.01 (0.96–1.05)	717 (19.5)	1096 (30.4)
2011	3924	128 659	3049.9 (2956.0–3146.9)	0.95 (0.91–1.00)	718 (18.3)	1109 (28.8)
2012	4377	137 062	3193.4 (3100.2–3289.5)	0.99 (0.95–1.04)	818 (18.7)	1184 (27.4)
2013	4652	136 877	3398.7 (3302.4–3497.8)	1.06 (1.01–1.11)*P for trend=0.0003	837 (18.0)	1220 (26.7)

*P values for likelihood ratio test.

Table 2. Median length of stay on ICU, median length of stay in hospital and associated costs for individuals with bronchiectasis

Bronchiectasis	Median length of stay on ICU (IQR) ^a	Median length of stay in hospital (IQR) ^a	Cost on NHS in terms of critical care admission (£)
Year			
2009	3.0 (1.0–5.0)	14 (7–24)	189 144
2010	3.5 (1.4–6.4)	14 (7–29)	232 596
2011	2.1 (1.0–5.1)	17 (7–33)	193 233
2012	3.8 (1.1–7.3)	16 (8–30)	501 828
2013	2.9 (1.4–7.3)	12 (6–25)	298 967

^aLength of stay in days.**Table 3.** Characteristics of individuals with bronchiectasis and COPD aged 70 or above admitted to ICU

	Individuals with bronchiectasis aged ≥70 years (n = 219)	Individuals with COPD aged ≥70 years (n = 8844)
Mean age at admission (SD)	76.6 (5.0)	76.9 (5.1)
Male (%)	98 (44.7)	4567 (51.6)
Mean APACHE II acute physiology score (SD)	13.3 (5.9)	12.5 (5.6)
Number of people intubated (%)	91 (41.6)	3727 (42.2)
Number of ICU deaths (%)	68 (31.1)	2042 (23.1)
Number of hospital deaths (%)	102 (46.6)	3215 (36.4)
Median length of stay on ICU (IQR)	2.9 (1.1–6.4)	2.8 (1.2–5.9)

Our study also demonstrates that both ICU mortality and in-hospital mortality were substantially higher in people with bronchiectasis compared with people with COPD, especially in those 70 years or above, despite similar acute physiological impairment and rates of intubation. On the basis of our findings, we estimate that the annual financial burden from bronchiectasis critical care admissions alone in the UK will be almost £500 000 by 2020, although this likely to be an underestimation,

as it assumes only single organ support and does not take into account inflation costs and changes to market forces.

We used a large, comprehensive dataset that includes admissions across almost all adult critical care centres in England, Wales and Northern Ireland, including specialist tertiary centres and district general hospitals. The ICNARC dataset has been shown to be broadly representative of the UK population of patients admitted to ICU.⁷ This has enabled us to estimate a realistic disease burden in terms of critical care admissions without sampling bias. The long study period has also enabled us to examine changes to age at admission, rates of intubation, ICU mortality, in-hospital mortality and length of stay.

One of the limitations of our study is that only summary data were available, hence we were unable to standardize our admission rates to account for the increase in the number of critical care units contributing to ICNARC. Hence, it is possible that the increase in admissions seen in our study is due, at least in part, to increased diagnosis of cases of bronchiectasis in the UK. Another potential limitation relates to the validity of and accuracy of diagnostic coding of ICU admission data. We have not validated the diagnosis of bronchiectasis or COPD within this dataset, but a previous study has shown that validity of admission coding within ICNARC is high.⁸ Furthermore, the demographic features of individuals with bronchiectasis in these data are consistent with the UK population of patients with bronchiectasis.^{9,10} However, given that the diagnosis of COPD is co-existent in 36% of individuals with bronchiectasis,¹¹ there is likely to be some diagnostic imprecision within our dataset, including undiagnosed Bronchiectasis–COPD overlap syndrome⁶ that may be included in the COPD population. Nevertheless, our study demonstrates a high ICU and in-hospital mortality rate in people with identified bronchiectasis, especially in those aged 70 years or above. Although this may be partly explained by other factors that influence mortality, this does not detract from our finding that survival in these individuals is poor.

There are sparse data on ICU admission rates in people with bronchiectasis, as well as outcomes with regards to mortality. A French study of 48 patients with bronchiectasis admitted to an ICU unit over a 10-year period suggested that age over 65 years and long-term oxygen therapy use were risk factors for increased mortality on ICU.¹² A single-centre study from 1996 to 2004 by Alzeer et al.¹³ reported that the mean age of patients

with bronchiectasis admitted to ICU was 63.5 years. This study also found that older patients with bronchiectasis admitted to ICU had a worse outcome in terms of survival.¹³ Both studies report variable ICU mortality rates, ranging from 19 to 34%,^{12,13} and long-term oxygen therapy was also associated with increased mortality.¹³

In summary, this first ever national study of ICU outcomes in patients with bronchiectasis in the UK has identified the increasing demand on critical care services and the financial impact of this on secondary care services. We have also highlighted the poor prognosis of individuals with bronchiectasis admitted to ICU, especially in the older population of 70 years or older. These data will aid updates in national bronchiectasis guidelines¹⁴ and help guide thresholds for admission to critical care.

Acknowledgements

The authors would like to thank Francesca Parrot from ICNARC for extracting the dataset and help with the summary analysis. We are grateful to the units submitting data to the ICNARC database. We would also like to thank BRONCH-UK, a Medical Research Council funded research collaboration in non-cystic fibrosis bronchiectasis (Reference MR/L011263/1).

Funding

Medical Research Council partnership in bronchiectasis, Bronch UK (www.bronch.ac.uk) (Reference MR/L011263/1). R.B.H. is funded by the GSK/BLF chair of Epidemiological Respiratory Research. V.N. is funded by the NIHR as an Academic Clinical Fellow.

Conflict of interest: None declared.

References

1. Ip M, Shum D, Lauder I, Lam WK, So SY. Effect of antibiotics on sputum inflammatory contents in acute exacerbations of bronchiectasis. *Respir Med* 1993; **87**:449–54.
2. Quint JK, Millett ER, Joshi M, Navaratnam V, Thomas SL, Hurst JR, et al. Changes in the incidence, prevalence and mortality of bronchiectasis in the UK from 2004 to 2013: a population-based cohort study. *The European respiratory journal*. 2015 Nov 5.
3. Navaratnam V, Millett E, Hurst J, Thomas S, Smeeth L, Hubbard R, et al. P81 the increasing secondary care burden of bronchiectasis in England. *Thorax* 2014; **69**:A111–2.
4. Roberts HJ, Hubbard R. Trends in bronchiectasis mortality in England and Wales. *Respir Med* 2010; **104**:981–5.
5. ICNARC. The Intensive Care National Audit and Research Centre. (www.icnarc.org)
6. Hurst JR, Elborn JS, De Soya A. COPD-bronchiectasis overlap syndrome. *Eur Respir J* 2015; **45**:310–3.
7. Harrison DA, Brady AR, Rowan K. Case mix, outcome and length of stay for admissions to adult, general critical care units in England, Wales and Northern Ireland: the Intensive Care National Audit & Research Centre Case Mix Programme Database. *Crit Care* 2004; **8**:R99–111.
8. de Keizer NF, Bonsel GJ, Goldfad C, Rowan KM. The added value that increasing levels of diagnostic information provide in prognostic models to estimate hospital mortality for adult intensive care patients. *Intensive Care Med* 2000; **26**:577–84.
9. Hill AT, Welham S, Reid K, Bucknall CE. British thoracic society national bronchiectasis audit 2010 and 2011. *Thorax* 2012; **67**:928–30.
10. Hill AT, Routh C, Welham S. National BTS bronchiectasis audit 2012: is the quality standard being adhered to in adult secondary care? *Thorax* 2013; **69**:292–4.
11. Navaratnam V, Millett E, Hurst JR, Thomas S, Smeeth L, Hubbard R, et al. *The Prevalence of Co-Existing Diseases Associated with the Aetiology of Bronchiectasis. C23 Environmental and Psychosocial Aspects of Asthma and COPD*. New York: American Thoracic Society, 2014; p. A4043–A.
12. Dupont M, Gacouin A, Lena H, Lavoue S, Brinchault G, Delaval P, et al. Survival of patients with bronchiectasis after the first ICU stay for respiratory failure. *Chest* 2004; **125**:1815–20.
13. Alzeer AH, Masood M, Basha SJ, Shaik SA. Survival of bronchiectatic patients with respiratory failure in ICU. *BMC Pulm Med* 2007; **7**:17.
14. Pasteur MC, Bilton D, Hill AT. British Thoracic Society guideline for non-CF bronchiectasis. *Thorax* 2010; **65**:i1–58.